arXiv:1406.2182v2 [math-ph] 21 Jan 2015

Elementary derivation of Weingarten functions of classical Lie groups

Marcel Novaes
Instituto de Física, Universidade Federal de Uberlândia, Uberlândia, MG, 38408-100, Brazil

Abstract

Integration of polynomials over the classical groups of unitary, orthogonal and symplectic matrices can be reduced to basic building blocks known as Weingarten functions. We present an elementary derivation of these functions.

1 Introduction

Integration with respect to matrix ensembles is the core of random matrix theory [1] and an important problem in many areas of mathematical physics. [2] Classical Lie groups of unitary, orthogonal and symplectic matrices, endowed with the corresponding Haar measure, constitute important classes of matrix ensembles. Integrals of functions which are polynomials in the matrix elements can be reduced to the general form

$$\int_G \prod_{k=1}^{n_1} U_{i_k j_k} \prod_{\ell=1}^{n_2} U^*_{m_\ell r_\ell} dU \equiv \left\langle \prod_{k=1}^{n} U_{i_k j_k} \prod_{\ell=1}^{n_2} U^*_{m_\ell r_\ell} \right\rangle_G , \tag{1.1}$$

where G denotes one of the groups and dU the corresponding Haar measure.

Initial investigations in the physics literature considered the unitary group $\mathcal{U}(N)$.[3, 4, 5, 6, 7, 8, 9] It was found that the basic integral can be written as a double sum over the symmetric group,

$$\left\langle \prod_{k=1}^{n} U_{i_k j_k} U^*_{m_k r_k} \right\rangle_{\mathcal{U}(N)} = \sum_{\tau,\sigma \in S_n} \mathrm{Wg}^U(\tau^{-1}\sigma) \prod_{k=1}^{n} [i_k = m_{\sigma(k)}][j_k = r_{\tau(k)}], \tag{1.2}$$

where $[i = m]$ is the same as $\delta_{i,m}$ and Wg^U is a function which depends only on the cycle structure of its argument. Independently of the physicists, Collins [10] rediscovered the problem and suggested Wg^U be called the Weingarten function of $\mathcal{U}(N)$.

The corresponding question for the orthogonal and symplectic groups has also been much studied. [11, 12, 13, 14] They too can be written as double sums over permutations, and the corresponding Weingarten functions Wg^O and Wg^{Sp} have been found. [15, 16] Later developments include new approaches (e.g. from Jucys-Murphy elements, [17, 18]) generalizations (e.g. to compact symmetric spaces [19, 20]) connections (e.g. to factorizations of permutations [21]) and applications (e.g. to different polynomial integrals, [22] to quantum mechanics [23, 24, 25]).

Previous works where Weingarten functions were obtained were based either on representation theory and Schur-Weyl duality,[10, 15] the theory of Gelfand pairs,[20] or Jucys-Murphy elements.[18] In contrast, we here derive Weingarten functions for the classical

compact groups by means of some elementary direct calculations (although we rely on some classical results that can, of course, be interpreted very naturally in the light of those theories).

The idea consists of five steps: 1) write the integrand as the derivative of a power sum function; 2) change basis from power sums to Schur functions; 3) perform the group integral; 4) revert back to power sums; 5) take the derivative to arrive at the result.

2 Required facts

Let us review some well known facts and establish notation and terminology. More detailed explanations of the concepts introduced below can be found in the classical monograph by MacDonald [26].

Everywhere, $\lambda \vdash n$ means λ is a partition of n, i.e. a weakly decreasing sequence of positive integers such that $\sum_i \lambda_i = n$. The number of non-zero parts is called the length of the partition and denoted by $\ell(\lambda)$. The partition with n unit parts is denoted 1^n. We define $2\lambda \equiv (2\lambda_1, 2\lambda_2, ...)$ and $\lambda \cup \lambda \equiv (\lambda_1, \lambda_1, \lambda_2, \lambda_2, ...)$.

2.1 Lie groups

The unitary group $\mathcal{U}(N)$ is the group of complex $N \times N$ matrices satisfying $U^\dagger U = 1$, where $U^\dagger$ is the transpose conjugate of U.

The orthogonal group $\mathcal{O}(N)$ is the subgroup of $\mathcal{U}(N)$ of real matrices. This is the isometry group in $\mathbb{R}^N$ with respect to the scalar product $\{u, v\} = \sum_{i=1}^N u_i v_i$.

Let $J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, where 0 and 1 are, respectively, the zero and the identity matrix in N dimensions. The unitary symplectic group $Sp(2N)$ is the subgroup of $\mathcal{U}(2N)$ of complex unitary matrices satisfying $U^D U = 1$, where $U^D = JU^T J^T$. If we define

$$\langle\langle u, v \rangle\rangle = \sum_{i=1}^{2N} u_i (Jv)_i = \sum_{i=1}^{N} u_i v_{i+N} - u_{i+N} v_i, \tag{2.1}$$

then U^D plays the role of adjoint of U, in the sense that $\langle\langle u, Uv \rangle\rangle = \langle\langle U^D u, v \rangle\rangle$.

2.2 Permutation groups

The group of permutations of n elements is S_n. The cycle type of $\pi \in S_n$ is the partition $\lambda \vdash n$ whose parts are the lengths of the cycles of π (the cycle type of the identity in S_n is 1^n). We denote by $\mathcal{C}_\lambda$ the conjugacy class of all permutations of cycle type λ. The length of a permutation is $\ell(\pi) = \ell(\lambda)$ if $\pi \in \mathcal{C}_\lambda$, and the sign is $s(\pi) = (-1)^{n-\ell(\pi)}$. We multiply permutations from right to left, e.g. $(13)(12) = (123)$.

The quantity $\chi_\lambda(\mu) = \chi_\lambda(\pi)$ is the character of the irreducible representation of S_n labeled by λ, calculated for a permutation $\pi \in \mathcal{C}_\mu$. The value $d_\lambda = \chi_\lambda(1^n)$ is the dimension of said representation. Characters satisfy the orthogonality relation

$$\sum_{\tau \in S_n} \chi_\mu(\tau) \chi_\lambda(\tau\sigma) = \delta_{\mu,\lambda} n! \frac{\chi_\lambda(\sigma)}{d_\lambda}. \tag{2.2}$$

The group S_{2n} has a subgroup called the hyperoctahedral, H_n, with $|H_n| = 2^n n!$ elements, which is the stabilizer of the permutation $(12)(34)\cdots(2n-1\ 2n)$. The coset $S_{2n}/H_n = \mathcal{M}_n$

can be represented by permutations called matchings: $\sigma \in \mathcal{M}_n$ if and only if $\sigma(2i-1) < \sigma(2i)$ and $\sigma(2i-1) < \sigma(2i+1)$.

Given a permutation $\tau \in S_{2n}$, let $\mathcal{G}_\tau$ be a graph with vertices labeled from 1 to $2n$ and edges of the forms $\{2i-1, 2i\}$ and $\{\tau(2i-1), \tau(2i)\}$, $1 \le i \le n$. Since each vertex belongs to one edge of each form, all connected components of $\mathcal{G}_\tau$ are cycles of even length. The coset type of τ is the partition of n whose parts are half the number of edges in the connected components of $\mathcal{G}_\tau$.

Two permutations τ, σ in S_{2n} have the same coset type if and only if they belong to the same double coset, i.e. if $\tau = h_1 \sigma h_2$ with $h_1, h_2, \in H_n$. We denote by $[\tau]$ the coset type of τ. We may denote by H_τ the double coset of τ. Its order is

$$|H_\tau| = \frac{4^n n! |\mathcal{C}_{[\tau]}|}{2^{\ell([\tau])}}. \tag{2.3}$$

Introduce a map from S_n to S_{2n}, taking a permutation $\pi \in \mathcal{C}_\lambda$ into another permutation $\widetilde{\pi} \in \mathcal{C}_{2\lambda}$. This map is such that the image of the cycle $(i_1, i_2, \cdots, i_r)$ in π is the cycle $(2i_1-1, 2i_1, 2i_2-1, 2i_2, \cdots, 2i_r-1, 2i_r)$ in $\widetilde{\pi}$. This map is useful because the coset type of $\widetilde{\pi}$ is λ.

The average

$$\omega_\lambda(\tau) = \frac{1}{|H_n|} \sum_{\xi \in H_n} \chi_{2\lambda}(\tau\xi) \tag{2.4}$$

is called a zonal spherical function. It is obviously invariant under (left and right) action of H_n and hence depends only on the coset type of its argument. These functions satisfy

$$\sum_{\tau \in \mathcal{M}_n} \omega_\lambda(\tau)\omega_\mu(\tau^{-1}\sigma) = \delta_{\lambda,\mu} \frac{(2n)!}{2^n n!} \frac{\omega_\lambda(\sigma)}{d_{2\lambda}}, \tag{2.5}$$

and also

$$\sum_{\lambda \vdash n} d_{2\lambda}\omega_\lambda(\sigma)\omega_\lambda(\tau) = \frac{(2n)!}{|H_\tau|} \delta_{[\sigma],[\tau]}. \tag{2.6}$$

The average

$$\psi_\lambda(\tau) = \frac{1}{|H_n|} \sum_{\xi \in H_n} \chi_{\lambda\cup\lambda}(\tau\xi) s(\xi), \tag{2.7}$$

where $s(\xi)$ is the sign of ξ, is called a twisted zonal spherical function. Under action by H_n it is invariant up to sign: $\psi_\lambda(\tau\xi) = s(\xi)\psi_\lambda(\tau)$. These functions satisfy

$$\sum_{\tau \in \mathcal{M}_n} \psi_\lambda(\tau)\psi_\mu(\tau^{-1}\sigma) = \delta_{\lambda,\mu} \frac{(2n)!}{2^n n!} \frac{\psi_\lambda(\sigma)}{d_{\lambda\cup\lambda}}, \tag{2.8}$$

and also

$$\sum_{\lambda \vdash n} d_{\lambda\cup\lambda}\psi_\lambda(\sigma)\psi_\lambda(\tau) = \frac{(2n)!}{|H_\sigma|} s(\sigma)s(\tau)\delta_{[\sigma],[\tau]}. \tag{2.9}$$

2.3 Power sum symmetric functions

Power sum symmetric functions of matrix argument are given by

$$p_\lambda(X) = \prod_{i=1}^{\ell(\lambda)} p_{\lambda_i}(X), \tag{2.10}$$

with $p_m(X) = \mathrm{Tr}(X^m)$. They are clearly symmetric functions of the eigenvalues of X. If X has dimension N and $\pi \in S_n$ has cycle type λ, then

$$p_\pi(X) = p_\lambda(X) = \sum_{i_1,\ldots,i_n=1}^{N} \prod_{k=1}^{n} X_{i_k i_{\pi(k)}}. \tag{2.11}$$

Power sums are produced from some summations. Let δ_{ab} denote the usual Kronecker delta function. Given $j = (j_1, j_2, \ldots j_n)$ and $m = (m_1, m_2, \ldots, m_n)$, define the function

$$\delta_\tau[j,m] = \prod_{k=1}^{n} \delta_{j_k m_{\tau(k)}}, \tag{2.12}$$

for any $\tau \in S_n$. Then we have

$$\sum_{j_1\cdots j_n=1}^{N} \sum_{m_1\cdots m_n=1}^{N} \delta_\tau[j,m]\delta_\sigma[j,m] \prod_{k=1}^{n} y_{j_k} y_{m_k} = p_{\tau^{-1}\sigma}(x), \tag{2.13}$$

where

$$x_i = y_i^2, \quad 1 \le i \le N. \tag{2.14}$$

Let $e(j)$ be the vector with elements given by $(e(j))_i = \delta_{ij}$. With the definitions from Section 2.1 we have

$$\{e(j_1), e(j_2)\} = \delta_{j_1,j_2}, \quad \langle\!\langle e(j_1), e(j_2) \rangle\!\rangle = \delta_{j_1+N,j_2} - \delta_{j_1,j_2+N}. \tag{2.15}$$

For any $\tau \in S_{2n}$, define

$$\Delta_\tau[j] = \prod_{k=1}^{n} \{e(j_{\tau(2k-1)}), e(j_{\tau(2k)})\}, \tag{2.16}$$

and

$$\Delta'_\tau[j] = \prod_{k=1}^{n} \langle\!\langle e(j_{\tau(2k-1)}), e(j_{\tau(2k)}) \rangle\!\rangle. \tag{2.17}$$

Notice that the behavior of these functions under action of the hyperoctahedral group is $\Delta_{\tau\xi} = \Delta_\tau$ and $\Delta'_{\tau\xi} = s(\xi)\Delta'_\tau$. The summations that produce power sums are

$$\sum_{j_1\cdots j_{2n}=1}^{N} \Delta_\tau[j]\Delta_\sigma[j] \prod_{k=1}^{2n} y_{j_k} = p_{[\tau^{-1}\sigma]}(x), \tag{2.18}$$

and

$$\sum_{j_1\cdots j_{2n}=1}^{2N} \Delta'_\tau[j]\Delta'_\sigma[j] \prod_{k=1}^{2n} y_{j_k} = (-1)^n s(\tau)s(\sigma)(-2)^{\ell([\tau^{-1}\sigma])} p_{[\tau^{-1}\sigma]}(x), \tag{2.19}$$

where $[\tau]$ is the coset type of τ, while the variables satisfy $y_{i+N} = y_i$ and (2.14).

2.4 Jack polynomials

Power sums and Schur functions are related by

$$s_\lambda(X) = \sum_{\pi\in S_n} \chi_\lambda(\pi) p_\pi(X) = \sum_{\mu\vdash n} |\mathcal{C}_\mu| \chi_\lambda(\mu) p_\mu(X), \tag{2.20}$$

and

$$p_\lambda(X) = \sum_{\mu\vdash n} \chi_\mu(\lambda) s_\mu(X). \tag{2.21}$$

Schur functions are generalized by Jack polynomials, $J_\lambda^{(\alpha)}(x)$, which depend on the parameter α. They are recovered when $\alpha = 1$:

$$J_\lambda^{(1)}(x) = \frac{n!}{d_\lambda} s_\lambda(x). \tag{2.22}$$

The polynomials $J_\lambda^{(2)}$ and $J_\lambda^{(1/2)}$ are called zonal polynomials and quaternion zonal polynomials, respectively. The former are related to power sums according to

$$J_\lambda^{(2)}(X) = \sum_{\mu \vdash n} |\mathcal{C}_\mu| 2^{n-\ell(\mu)} \omega_\lambda(\mu) p_\mu(X), \tag{2.23}$$

and

$$p_{[\tau]}(x) = \frac{2^n n!}{(2n)!} \sum_{\lambda \vdash n} d_{2\lambda} \omega_\lambda(\tau) J_\lambda^{(2)}(x). \tag{2.24}$$

For the latter, the first relation is

$$J_\lambda^{(1/2)}(X) = (-1)^n \sum_{\mu \vdash n} |\mathcal{C}_\mu| \psi_\lambda(\widetilde{\pi}) p_\mu(X), \tag{2.25}$$

where π is any permutation of cycle type μ and $\widetilde{\pi}$ is its image under the map introduced in Section 2.2. The second relation is

$$p_{[\tau]}(x) = \frac{(-4)^n n!}{(2n)!} \left(\frac{-1}{2}\right)^{\ell([\tau])} \sum_{\lambda \vdash n} d_{\lambda \cup \lambda} s(\tau) \psi_\lambda(\tau) J_\lambda^{(1/2)}(x), \tag{2.26}$$

valid for any $\tau \in S_{2n}$.

The value of the Jack polynomial when all arguments are equal to 1 is given by

$$J_\lambda^{(\alpha)}(1^N) = \alpha^n \prod_{j=1}^{\ell(\lambda)} \frac{\Gamma(\lambda_j + (N-j+1)/\alpha)}{\Gamma((N-j+1)/\alpha)}. \tag{2.27}$$

Jack polynomials arise from integrals of Schur functions over Lie groups. Let dU, dO and dS denote the normalized Haar measures over $\mathcal{U}(N)$, $\mathcal{O}(N)$ and $Sp(2N)$, respectively. The orthogonality relation

$$\int_{\mathcal{U}(N)} dU s_\lambda(A^\dagger U) s_\mu(B U^\dagger) = \delta_{\mu,\lambda} \frac{J_\lambda^{(1)}(A^\dagger B)}{J_\lambda^{(1)}(1^N)} \tag{2.28}$$

holds because Schur functions are actually irreducible characters of the unitary group. We also have the relations

$$\int_{\mathcal{O}(N)} dO s_\mu(AO) = \delta_{\mu,2\lambda} \frac{J_\lambda^{(2)}(A^T A)}{J_\lambda^{(2)}(1^N)}, \tag{2.29}$$

and

$$\int_{Sp(2N)} dS s_\mu(AS) = \delta_{\mu,\lambda\cup\lambda} 2^{-\ell(\lambda)} \frac{J_\lambda^{(1/2)}(A^D A)}{J_\lambda^{(1/2)}(1^N)}. \tag{2.30}$$

In the last equation A and S are $2N \times 2N$ complex matrices. All the above integrals vanish unless $\ell(\lambda) \leq N$.

3 Derivation

3.1 Unitary Group

We start from the basic identity

$$\prod_{k=1}^{n} U_{a_k b_k} = \frac{1}{n!} \prod_{k=1}^{n} \frac{\partial}{\partial A^*_{a_k b_k}} p_{1^n}(A^\dagger U), \tag{3.1}$$

which can be easily verified. We express power sums in terms of Schur functions, use (2.28) to perform the group integral, and then revert back to power sums to get

$$\int_{\mathcal{U}(N)} dU \prod_{k=1}^{n} U_{a_k b_k} U^*_{d_k c_k} = \frac{1}{n!^2} \sum_{\substack{\lambda \vdash n \\ \ell(\lambda) \le N}} \frac{d_\lambda}{J^{(1)}_\lambda(1^N)} \sum_{\pi \in S_n} \chi_\lambda(\pi) \prod_{k=1}^{n} \frac{\partial}{\partial A^*_{a_k b_k}} \frac{\partial}{\partial B_{m_k r_k}} p_\pi(A^\dagger B). \tag{3.2}$$

In order to take the derivative, we expand p_π as a trace,

$$p_\pi(A^\dagger B) = \sum_{i_1,\ldots,i_n=1}^{N} \sum_{j_1,\ldots,j_n=1}^{N} A^\dagger_{i_k j_k} B_{j_k i_{\pi(k)}}. \tag{3.3}$$

Taking the derivative and summing over the i's and j's produces

$$\sum_{\theta,\rho \in S_n} \prod_{k=1}^{n} \delta_{a_k, d_{\rho\theta^{-1}(k)}} \delta_{b_k, c_{\rho\pi^{-1}\theta^{-1}(k)}}. \tag{3.4}$$

Finally, we change variables as $\rho = \sigma\theta$ and then $\tau = \sigma\theta\pi^{-1}\theta^{-1}$. This last change preserves cycle type, so that $\chi_\lambda(\pi) = \chi_\lambda(\sigma^{-1}\tau) = \chi_\lambda(\tau^{-1}\sigma)$. The sum over π is then just $n!$, and we arrive at

$$\int_{\mathcal{U}(N)} dU \prod_{k=1}^{n} U_{a_k b_k} U^*_{d_k c_k} = \sum_{\sigma\tau \in S_n} \mathrm{Wg}^U(\tau^{-1}\sigma) \delta_\sigma[ad] \delta_\tau[bc], \tag{3.5}$$

where

$$\mathrm{Wg}^U(\tau^{-1}\sigma) = \frac{1}{n!} \sum_{\substack{\lambda \vdash n \\ \ell(\lambda) \le N}} \frac{d_\lambda}{J^{(1)}_\lambda(1^N)} \chi_\lambda(\tau^{-1}\sigma), \tag{3.6}$$

is the unitary Weingarten function.

3.2 Orthogonal Group

In the same spirit, we start by writing

$$\prod_{k=1}^{2n} O_{a_k b_k} = \frac{1}{(2n)!} \prod_{k=1}^{2n} \frac{\partial}{\partial A_{a_k b_k}} p_{1^{2n}}(A^T O). \tag{3.7}$$

Changing to Schur functions and using (2.29) we get

$$\frac{1}{(2n)!} \sum_{\substack{\lambda \vdash n \\ \ell(\lambda) \le N}} \frac{d_{2\lambda}}{J^{(2)}_\lambda(1^N)} \prod_{k=1}^{2n} \frac{\partial}{\partial A_{a_k b_k}} J^{(2)}_\lambda(A^T A). \tag{3.8}$$

We return to power sums in order to take the derivative. Taking $\pi \in S_n$ to be any permutation with cycle type μ, we write

$$\int_{\mathcal{O}(N)} dO \prod_{k=1}^{2n} O_{a_k b_k} = \frac{1}{(2n)!} \sum_{\substack{\lambda \vdash n \\ \ell(\lambda) \leq N}} \frac{d_{2\lambda}}{J_\lambda^{(2)}(1^N)} \sum_{\mu \vdash n} |\mathcal{C}_\mu| 2^{n-\ell(\mu)} \omega_\lambda(\mu) \prod_{k=1}^{2n} \frac{\partial}{\partial A_{a_k b_k}} p_\pi(A^T A). \quad (3.9)$$

The derivative

$$\prod_{k=1}^{2n} \frac{\partial}{\partial A_{a_k b_k}} p_\pi(A^T A) = \sum_{i_1,\ldots,i_n=1}^{N} \sum_{j_1,\ldots,j_n=1}^{N} \prod_{r=1}^{n} \frac{\partial}{\partial A_{a_{(2r-1)} b_{(2r-1)}}} \frac{\partial}{\partial A_{a_{(2r)} b_{(2r)}}} A^T_{i_r j_r} A_{j_r i_{\pi(r)}}, \quad (3.10)$$

will, upon summing over the i's and j's, always lead to a matching among the a's and a matching among the b's, i.e. something like $\Delta_\tau[a]\Delta_\sigma[b]$. By construction, these matchings will satisfy $[\tau^{-1}\sigma] = \mu$, and each such pair will appear exactly $2^{\ell(\mu)} n!/|\mathcal{C}_\mu|$ times for any π. Hence, we can write

$$\int_{\mathcal{O}(N)} dO \prod_{k=1}^{2n} O_{a_k b_k} = \sum_{\tau,\sigma \in \mathcal{M}_n} \mathrm{Wg}^O(\tau^{-1}\sigma) \Delta_\tau(a) \Delta_\sigma(b), \quad (3.11)$$

where

$$\mathrm{Wg}^O(\tau^{-1}\sigma) = \frac{2^n n!}{(2n)!} \sum_{\substack{\lambda \vdash n \\ \ell(\lambda) \leq N}} \frac{d_{2\lambda}}{J_\lambda^{(2)}(1^N)} \omega_\lambda(\tau^{-1}\sigma) \quad (3.12)$$

is the Weingarten function of $\mathcal{O}(N)$.

3.3 Symplectic Group

Let $\pi \in S_n$ be any permutation of cycle type μ and $\widetilde{\pi}$ be its image under the map in Section 2.2, so that $[\widetilde{\pi}] = \mu$. Calculations analogous to those in the previous section lead to

$$\int_{Sp(2N)} dS \prod_{k=1}^{2n} S_{a_k b_k} = \frac{(-1)^n}{(2n)!} \sum_{\substack{\lambda \vdash n \\ \ell(\lambda) \leq N}} \frac{d_{\lambda \cup \lambda}}{J_\lambda^{(1/2)}(1^N)} \sum_{\mu \vdash n} |\mathcal{C}_\mu| 2^{-\ell(\mu)} \psi_\lambda(\widetilde{\pi}) \prod_{k=1}^{2n} \frac{\partial}{\partial A_{a_k b_k}} p_\pi(A^D A), \quad (3.13)$$

where we have used (2.25) and (2.30). The derivative $\prod_{k=1}^{2n} \frac{\partial}{\partial A_{a_k b_k}} p_\pi(A^D A)$ leads to a matching among the a's and a matching among the b's, but with such minus signs as to produce $(-1)^n \Delta'_\tau[a]\Delta'_\sigma[b]$. Again, these matchings satisfy $[\tau^{-1}\sigma] = \mu$, and each such pair appears $2^{\ell(\mu)} n!/|\mathcal{C}_\mu|$ times for any π. Therefore,

$$\int_{Sp(2N)} dS \prod_{k=1}^{2n} S_{a_k b_k} = \sum_{\tau,\sigma \in \mathcal{M}_n} \mathrm{Wg}^{Sp}(\tau^{-1}\sigma) \Delta'_\tau[a] \Delta'_\sigma[b], \quad (3.14)$$

where

$$\mathrm{Wg}^{Sp}(\tau^{-1}\sigma) = \frac{n!}{(2n)!} \sum_{\substack{\lambda \vdash n \\ \ell(\lambda) \leq N}} \frac{d_{\lambda \cup \lambda}}{J_\lambda^{(1/2)}(1^N)} \psi_\lambda(\tau^{-1}\sigma) \quad (3.15)$$

is the Weingarten function of $Sp(2N)$.

Acknowledgments

Financial support from CNPq is gratefully acknowledged, as well as from grant 2012/00699-1, São Paulo Research Foundation (FAPESP).

References

[1] M.L. Mehta, *Random Matrices* (Academic Press, 2004).

[2] G. Akemann, J. Baik, P. Di Francesco (Ed.), *The Oxford handbook of random matrix theory* (Oxford University Press, 2011).

[3] D. Weingarten, J. Math. Phys. **19**, 999 (1978).

[4] M. Creutz, J. Math. Phys. **19**, 2043 (1978).

[5] S. Samuel, J. Math. Phys. **21**, 2695 (1980).

[6] P. A. Mello, J. Phys. A **23**, 4061 (1990).

[7] P.W. Brouwer and C.W.J. Beenakker, J. Math. Phys. **37**, 4904 (1996).

[8] S. Aubert and C. S. Lam, J. Math. Phys. **44**, 6112 (2003).

[9] M. Degli Esposti and A. Knauf, J. Math. Phys. **45**, 4957 (2004).

[10] B. Collins, Int. Math. Res. Not. **17**, 953 (2003).

[11] T. Gorin, J. Math. Phys. **43**, 3342 (2002).

[12] T. Prosen, T.H. Seligman and H. A. Weidenmüller, J. Math. Phys. **43**, 5135 (2002).

[13] T. Gorin and G. V. López, J. Math. Phys. **49**, 013503 (2008).

[14] D. Braun, J. Phys. A: Math. Gen. **39**, 14581 (2006).

[15] B. Collins and P. Śniady, Comm. Math. Phys. **264**, 773 (2006).

[16] B. Collins and S. Matsumoto, J. Math. Phys. **50**, 113516 (2009).

[17] J. Novak, Banach Center Publ. **89**, 231 (2010).

[18] P. Zinn-Justin, Lett. Math. Phys. **91**, 119 (2010).

[19] B. Collins and M. Stolz, Ann. Prob. **36**, 876 (2008).

[20] S. Matsumoto, Random Matrices: Theory Appl. **2**, 1350001 (2013).

[21] S. Matsumoto and J. Novak, Int. Math. Res. Not. 2013(2), 362 (2013).

[22] T. Banica, B. Collins and J.-M. Schlenker, J. Combinat. Theory A **118**, 778 (2011).

[23] M. Žnidarič, C. Pineda and I. García-Mata, Phys. Rev. Lett. **107**, 080404 (2011).

[24] M. Novaes, J. Phys. A: Math. Theor. **46**, 502002 (2013).

[25] G. Berkolaiko and J. Kuipers, J. Math. Phys. **54**, 112103 (2013).

[26] I.G. Macdonald, *Symmetric Functions and Hall Polynomials*, 2nd edn. (Oxford University Press, Oxford, 1995).